

Coincidence Subwavelength Interference by a Classical Thermal Light

Kaige Wang^{1,2} and De-zhong Cao²

1.CCAST (World Laboratory), P. O. Box 8730, Beijing 100080

2.Department of Physics, Applied Optics Beijing Area Major Laboratory,
Beijing Normal University, Beijing 100875, China

We show that a thermal light random in transverse direction can perform subwavelength double slit interference in a joint-intensity measurement. This is the classical version of quantum lithography, and it can be explained with the correlation of rays instead of the entanglement of photons.

PACS number(s): 42.50.Dv, 42.50.St, 42.25.Hz

Noise is usually considered as harmful to communication and interference. Disorder and chaotic fluctuation may fade information and interference patterns. Can a noise be used as a signal carrier in communication and a source in interference? This study is aroused from a recent debate in quantum optics. In the last decade, the theoretical and experimental studies have shown that an entangled photon pair generated by the spontaneous parametric down-conversion (SPDC) exhibits some peculiar effects, such as sub-wavelength lithography, [1]- [8] coincidence or "ghost" imaging and interference. [9]- [16] Indeed, these effects had not been realized in the classical optics until that time. Especially, the sub-wavelength interference effect was ever one of the paradoxes relating to violations of the quantum mechanical uncertainty principle. Recently, Bennink et al have shown in their experiment that the coincidence imaging and the coincidence interference can be carried out by a classical light source. [17] Therefore, this demonstrates that the classical correlation plays the same (or similar) role as the quantum entanglement. The recent theoretical analysis [18] has shown that a thermal or quasi-thermal source can be charged with such a classical correlation. On the other hand, the sub-wavelength lithography is considered as a nonclassical interference surpassing the Rayleigh diffraction limit. Physically, this effect is explained with the quantum entanglement of photons [3] [4] or the photonic de Broglie wavelength of a multiphoton wavepacket. [1] [2] [5] A similar question has emerged: can the sub-wavelength interference be carried out by a classical beam, too? In this work, we demonstrate that a thermal light random in the transverse plane possesses a similar second-order spatial correlation to that of a photon pair perfectly entangled in the transverse wavevector. The difference between them is that the former has the correlation of the wavevector-degeneracy and the latter has the correlation of the wavevector-conservation. Therefore, a thermal light can perform not only the ghost imaging and interference as shown in the experiments, but also the sub-wavelength interference. Due to the different way in correlation, the observation of the sub-wavelength interference pattern is taken by a coincidence detection at a pair of symmetric positions. The result can be used to explain two kinds of macroscopic observation of sub-wavelength interference occurring in the SPDC of type I crystal. [19]

We consider the first-order and second-order correlations of the fields for a quantum system consisting of a photon pair and a classical thermal light. The beam propagates with a central wavevector $\mathbf{k}_0$ and frequency ω_0 . For a general two-photon state, the quantum wavefunction in the Schrödinger picture is written as $|\Psi\rangle = \int d\mathbf{q}_1 d\mathbf{q}_2 d\omega_1 d\omega_2 C(\mathbf{q}_1, \omega_1; \mathbf{q}_2, \omega_2) a_1^\dagger(\mathbf{q}_1, \omega_1) a_2^\dagger(\mathbf{q}_2, \omega_2) |0\rangle$, where $\mathbf{q}_i$ and ω_i ($i = 1, 2$) are the transverse wavevector (or the spatial frequency) and the frequency deviation from the central frequency, respectively. We assume that two photons are distinguishable by their polarizations. By taking into account the commutation relation of the field operators, we obtain the first- and second-order spectral correlations for the two-photon state $|\Psi\rangle$

$$\langle a_1^\dagger(\mathbf{q}, \omega) a_1(\mathbf{q}', \omega') \rangle = \int d\mathbf{q}_2 d\omega_2 C^*(\mathbf{q}, \omega; \mathbf{q}_2, \omega_2) C(\mathbf{q}', \omega'; \mathbf{q}_2, \omega_2), \quad (1a)$$

$$\langle a_2^\dagger(\mathbf{q}, \omega) a_2(\mathbf{q}', \omega') \rangle = \int d\mathbf{q}_1 d\omega_1 C^*(\mathbf{q}_1, \omega_1; \mathbf{q}, \omega) C(\mathbf{q}_1, \omega_1; \mathbf{q}', \omega'), \quad (1b)$$

and

$$\langle a_1^\dagger(\mathbf{q}_1, \omega_1) a_2^\dagger(\mathbf{q}_2, \omega_2) a_2(\mathbf{q}_2', \omega_2') a_1(\mathbf{q}_1', \omega_1') \rangle = C^*(\mathbf{q}_1, \omega_1; \mathbf{q}_2, \omega_2) C(\mathbf{q}_1', \omega_1'; \mathbf{q}_2', \omega_2'), \quad (2)$$

respectively. Equation (2) shows explicitly separable product of the wavefunction and its conjugate one, reflecting the feature of quantum wavepacket. This implies that the wavevector and frequency correlation may exist only within the same (positive or negative) frequency component. For the perfectly entangled two-photon state generated in SPDC, it has $C(\mathbf{q}_1, \omega_1; \mathbf{q}_2, \omega_2) = \delta(\mathbf{q}_1 + \mathbf{q}_2) \delta(\omega_1 + \omega_2)$ owing to the conservation of momentum and energy in the SPDC process. Therefore, one obtains

$$\langle a_i^\dagger(\mathbf{q}, \omega) a_i(\mathbf{q}', \omega') \rangle = \delta(\mathbf{q} - \mathbf{q}') \delta(\omega - \omega'), \quad (i = 1, 2) \quad (3)$$

$$\langle a_1^\dagger(\mathbf{q}_1, \omega_1) a_2^\dagger(\mathbf{q}_2, \omega_2) a_2(\mathbf{q}_2', \omega_2') a_1(\mathbf{q}_1', \omega_1') \rangle = \delta(\mathbf{q}_1 + \mathbf{q}_2) \delta(\omega_1 + \omega_2) \delta(\mathbf{q}_1' + \mathbf{q}_2') \delta(\omega_1' + \omega_2'). \quad (4)$$

The results are also true for two photons with the same polarizations.

Then we consider a classical thermal light. We assume a monochromatic plane wave $E_0 \exp[i(k_0 z - \omega_0 t)]$ illuminating a scattering material containing disorder scattering centers. After scattering, the field is written as $E(\mathbf{x}, z, t) = \int E(\mathbf{q}) \exp[i(\mathbf{q} \cdot \mathbf{x} + k_z z - \omega_0 t)] d\mathbf{q}$ where $\mathbf{q}$ is the transverse wavevector introduced by the random scattering and satisfies $|\mathbf{q}|^2 + k_z^2 = k_0^2$. Hence, $E(\mathbf{q})$ is the stochastic variable obeying the Gaussian statistics. However, the scattering waves with different transverse wavevectors are independent statistically. If $|\mathbf{q}| \ll k_0$, the scattering field is approximately written as $E(\mathbf{x}, z, t) = A(\mathbf{x}) \exp[i(k_0 z - \omega_0 t)]$ where $A(\mathbf{x}) = \int E(\mathbf{q}) \exp[i\mathbf{q} \cdot \mathbf{x}] d\mathbf{q}$ is the slowly varying envelope. As a result, we have defined a monochromatic thermal light random in both strength and propagation direction. According to Wiener-Khinchine theorem, the first-order spectral correlation must satisfy

$$\langle E^*(\mathbf{q}) E(\mathbf{q}') \rangle = S(\mathbf{q}) \delta(\mathbf{q} - \mathbf{q}'), \quad (5)$$

where $S(\mathbf{q})$ is the power spectrum of the spatial frequency. Obviously, Eqs. (3) and (5) indicate the same incoherence of the first-order for an ideal two-photon entangled state and a thermal light with a wide bandwidth. Though the beam is monochromatic, the disorder in spatial frequency washes out the interference pattern.

For any thermal statistics, the high-order correlation can be represented by the first-order ones. Hence, the second-order spectral correlation of the thermal light is written as

$$\begin{aligned} & \langle E^*(\mathbf{q}_1) E^*(\mathbf{q}_2) E(\mathbf{q}'_2) E(\mathbf{q}'_1) \rangle \\ &= \langle E^*(\mathbf{q}_1) E(\mathbf{q}'_1) \rangle \langle E^*(\mathbf{q}_2) E(\mathbf{q}'_2) \rangle + \langle E^*(\mathbf{q}_1) E(\mathbf{q}'_2) \rangle \langle E^*(\mathbf{q}_2) E(\mathbf{q}'_1) \rangle \\ &= S(\mathbf{q}_1) S(\mathbf{q}_2) [\delta(\mathbf{q}_1 - \mathbf{q}'_1) \delta(\mathbf{q}_2 - \mathbf{q}'_2) + \delta(\mathbf{q}_1 - \mathbf{q}'_2) \delta(\mathbf{q}_2 - \mathbf{q}'_1)]. \end{aligned} \quad (6)$$

Different from the case of the two-photon entanglement, in which two photons with the opposite wavevectors are correlated, Eq. (6) shows that two rays with the degenerate wavevectors are correlated. When the thermal light is split into two beams in the two arms of a beam splitter, the correlated degenerate rays are spatially separated. This is the origin of the coincidence imaging and coincidence interference for a classical source. [17] As a matter of fact, in any coherent beam, there is no such a correlation between rays, so the coincidence imaging can not occur.

Now, we discuss the double slit interference in a joint-intensity observation. Let a beam illuminate a double slit with the slit-width b and the distance between two slits d . The double slit and the detection plane are placed at the two focal planes of a lens with the focal length f . The Fourier transform of the double-slit function is written as $\tilde{T}(q) = (2b/\sqrt{2\pi}) \text{sinc}(qb/2) \cos(qd/2)$. For simplicity, we omit the time variable. This can be done by using a narrow frequency filter in beam. According to Ref. [19], the second-order spatial correlation of the field in the detection plane is obtained to be

$$\begin{aligned} & \langle E_d^*(x_1) E_d^*(x_2) E_d(x_2) E_d(x_1) \rangle \\ &= \frac{k_0^2}{(2\pi f)^2} \int \tilde{T}(\frac{k_0 x_1}{f} - q_1) \tilde{T}(\frac{k_0 x_2}{f} - q_2) \tilde{T}(\frac{k_0 x_2}{f} - q'_2) \tilde{T}(\frac{k_0 x_1}{f} - q'_1) \\ & \quad \times \langle E^*(q_1) E^*(q_2) E(q'_2) E(q'_1) \rangle dq_1 dq_2 dq'_2 dq'_1. \end{aligned} \quad (7)$$

For an ideal two-photon entangled state, by using Eq. (4), one has $\langle E_d^*(x_1) E_d^*(x_2) E_d(x_2) E_d(x_1) \rangle \propto \tilde{T}^2[\frac{k_0}{f}(x_1 + x_2)]$. If one uses a two-photon detector to scan the position, the sub-wavelength interference fringe with the perfect visibility can be observed. Since the two-photon detector is not available at present, in the experiment, the two-photon detection is carried out by a coincidence measurement (CM) of two orthogonally polarized photons. [2] [4] However, for a thermal light described above, one obtains

$$\begin{aligned} & \langle E_d^*(x_1) E_d^*(x_2) E_d(x_2) E_d(x_1) \rangle \\ &= \frac{k_0^2}{(2\pi f)^2} \left\{ \int \tilde{T}^2(\frac{k_0 x_1}{f} - q) S(q) dq \int \tilde{T}^2(\frac{k_0 x_2}{f} - q) S(q) dq + \left[\int \tilde{T}(\frac{k_0 x_1}{f} - q) \tilde{T}(\frac{k_0 x_2}{f} - q) S(q) dq \right]^2 \right\}. \end{aligned} \quad (8)$$

In the broadband limit, we set $S(q) \approx S(0)$ and obtain approximately

$$\langle E_d^*(x_1) E_d^*(x_2) E_d(x_2) E_d(x_1) \rangle = \frac{k_0^2 S^2(0)}{2\pi f^2} \left\{ \tilde{T}^2(0) + \tilde{T}^2[\frac{k_0}{f}(x_1 - x_2)] \right\}. \quad (9)$$

When two detectors are placed in a pair of symmetric positions, $x_1 = -x_2 = x$, to perform a joint-intensity measurement, one observes the sub-wavelength interference fringe with the 50% visibility. In the general case, we set the Gaussian spectrum $S(q) = (\sqrt{2\pi}w)^{-1} \exp[-q^2/(2w^2)]$. Figure 1 shows the interference fringe as function of the

normalized bandwidth $W = wb/(2\pi)$. At very small bandwidth, the interference fringe is "normal", i.e. the same as that for a coherent beam. As the bandwidth is increasing, the sub-wavelength interference effect is appearing and growing.

Physically, the difference of the sub-wavelength interference effects for the two-photon entangled state and the thermal light comes from the manners of the correlation, that is, the wavevector conservation and the wavevector degeneracy. In Fig. 2, we illustrate the mechanism of the sub-wavelength interference effect of the classical thermal correlation. Due to the correlation of wavevector degeneracy, a pair of correlated rays $E(q)$ and $E^*(q)$ interfere with another pair of correlated rays $E(q')$ and $E^*(q')$, resulting in twice the optical path difference.

The beam generated in SPDC may incorporate both the quantum entanglement and the classical thermal correlation. A plane-wave pump field activates a $\chi^{(2)}$ nonlinear crystal, the basic unitary transformation is described by [16]

$$a_m(\mathbf{q}, \omega) = U_m(\mathbf{q}, \omega)a_m^{in}(\mathbf{q}, \omega) + V_m(\mathbf{q}, \omega)a_n^{in\dagger}(-\mathbf{q}, -\omega) \quad (m \neq n = s, i), \quad (10)$$

where $a_m(\mathbf{q}, \omega)$ and $a_m^{in}(\mathbf{q}, \omega)$ are the output and input field operators, respectively. The first-order correlation is obtained to be

$$\langle a_m^\dagger(\mathbf{q}, \omega)a_m(\mathbf{q}', \omega') \rangle = |V_m(\mathbf{q}, \omega)|^2 \delta(\mathbf{q} - \mathbf{q}') \delta(\omega - \omega'). \quad (11)$$

For type I crystal, one may omit the subscript in Eq. (10) and the second-order correlation is written as

$$\begin{aligned} & \langle a^\dagger(\mathbf{q}_1, \omega_1)a^\dagger(\mathbf{q}_2, \omega_2)a(\mathbf{q}'_2, \omega'_2)a(\mathbf{q}'_1, \omega'_1) \rangle \\ &= V^*(\mathbf{q}_1, \omega_1)V(\mathbf{q}'_1, \omega'_1)U^*(-\mathbf{q}_1, -\omega_1)U(-\mathbf{q}'_1, -\omega'_1)\delta(\mathbf{q}_1 + \mathbf{q}_2)\delta(\omega_1 + \omega_2)\delta(\mathbf{q}'_1 + \mathbf{q}'_2)\delta(\omega'_1 + \omega'_2) \\ &+ |V(\mathbf{q}_1, \omega_1)V(\mathbf{q}_2, \omega_2)|^2[\delta(\mathbf{q}_1 - \mathbf{q}'_1)\delta(\mathbf{q}_2 - \mathbf{q}'_2)\delta(\omega_1 - \omega'_1)\delta(\omega_2 - \omega'_2) + \delta(\mathbf{q}_1 - \mathbf{q}'_2)\delta(\mathbf{q}_2 - \mathbf{q}'_1)\delta(\omega_1 - \omega'_2)\delta(\omega_2 - \omega'_1)]. \end{aligned} \quad (12)$$

The first term shows the same correlation as the two-photon entangled state, whereas the second term shows the same correlation as the thermal light. Note that this result is valid for general coupling gain of SPDC, even if the converted beam contains a large number of photons. When the gain is very small, the second term is negligible in comparison with the first term and the converted field is approximately in a two-photon entangled state. As the gain is increasing, two kinds of macroscopic observation of the sub-wavelength interference co-exist. [19] For type II crystal, however, the second-order correlation of two orthogonally polarized beams is obtained to be

$$\begin{aligned} & \langle a_m^\dagger(\mathbf{q}_1, \omega_1)a_n^\dagger(\mathbf{q}_2, \omega_2)a_n(\mathbf{q}'_2, \omega'_2)a_m(\mathbf{q}'_1, \omega'_1) \rangle \quad (m \neq n = s, i) \\ &= V_m^*(\mathbf{q}_1, \omega_1)V_m(\mathbf{q}'_1, \omega'_1)U_n^*(-\mathbf{q}_1, -\omega_1)U_n(-\mathbf{q}'_1, -\omega'_1)\delta(\mathbf{q}_1 + \mathbf{q}_2)\delta(\omega_1 + \omega_2)\delta(\mathbf{q}'_1 + \mathbf{q}'_2)\delta(\omega'_1 + \omega'_2) \\ &+ |V_m(\mathbf{q}_1, \omega_1)V_n(\mathbf{q}_2, \omega_2)|^2\delta(\mathbf{q}_1 - \mathbf{q}'_1)\delta(\mathbf{q}_2 - \mathbf{q}'_2)\delta(\omega_1 - \omega'_1)\delta(\omega_2 - \omega'_2). \end{aligned} \quad (13)$$

The first term is the same as that for type I case, but the second term does not represent a thermal correlation. One can observe the quantum subwavelength interference only. [19] The thermal correlation in type-II crystal can be recovered as long as the signal or idler beam is extracted by a polarization beamsplitter. The second-order correlation for the signal/idler beam reads

$$\begin{aligned} & \langle a_m^\dagger(\mathbf{q}_1, \omega_1)a_m^\dagger(\mathbf{q}_2, \omega_2)a_m(\mathbf{q}'_2, \omega'_2)a_m(\mathbf{q}'_1, \omega'_1) \rangle \quad (m = s, i) \\ &= |V_m(\mathbf{q}_1, \omega_1)V_n(\mathbf{q}_2, \omega_2)|^2[\delta(\mathbf{q}_1 - \mathbf{q}'_1)\delta(\mathbf{q}_2 - \mathbf{q}'_2)\delta(\omega_1 - \omega'_1)\delta(\omega_2 - \omega'_2) + \delta(\mathbf{q}_1 - \mathbf{q}'_2)\delta(\mathbf{q}_2 - \mathbf{q}'_1)\delta(\omega_1 - \omega'_2)\delta(\omega_2 - \omega'_1)]. \end{aligned} \quad (14)$$

We obtain the thermal correlation for either signal or idler beam.

In conclusion, we show a classical version of quantum subwavelength lithography. This demonstrates that the classical correlation of a thermal light plays the similar role as the quantum entanglement of two photons in the optical subwavelength interference, in addition to the coincidence imaging and interference.

This research was supported by the National Fundamental Research Program of China with No. 2001CB309310, and the National Natural Science Foundation of China, Project Nos. 60278021 and 10074008.


-
- [1] J. Jacobson, G. Björk, I. Chuang, and Y. Yamamoto, Phys. Rev. Lett. **74**, 4835 (1995).
 - [2] E. J. S. Fonseca, C. H. Monken, and S. Pádua, Phys. Rev. Lett. **82**, 2868 (1999).
 - [3] A. N. Boto, P. Kok, D. S. Abrams, S. L. Braunstein, C. P. Williams, and J. P. Dowling, Phys. Rev. Lett. **85**, 2733 (2000).
 - [4] M. D'Angelo, M. V. Chekhova, and Y. Shih, Phys. Rev. Lett. **87**, 013602 (2001).

- [5] E. J. S. Fonseca, Z. Paulini, P. Nussenzveig, C. H. Monken, and S. Pádua, Phys. Rev. A **63**, 043819 (2001).
- [6] E. M. Nagasako, S. J. Bentley, R. W. Boyd, and G. S. Agarwal, Phys. Rev. A **64**, 043802 (2001).
- [7] K. Edamatsu, R. Shimizu, and T. Itoh, Phys. Rev. Lett. **89**, 213601 (2002).
- [8] R. Shimizu, K. Edamatsu, and T. Itoh, Phys. Rev. A **67**, 041805(R) (2003).
- [9] D. V. Strekalov, A. V. Sergienko, D. N. Klyshko, and Y. H. Shih, Phys. Rev. Lett. **74**, 3600 (1995); T. B. Pittman, Y. H. Shih, D. V. Strekalov, and A. V. Sergienko, Phys. Rev. A **52**, R3429 (1995).
- [10] P. H. S. Ribeiro, S. Pádua, J. C. Machado da Silva, and G. A. Barbosa, Phys. Rev. A **49**, 4176 (1994).
- [11] P. H. Souto Ribeiro and G. A. Barbosa, Phys. Rev. A **54**, 3489 (1996).
- [12] G. A. Barbosa, Phys. Rev. A **54**, 4473 (1996).
- [13] E. J. S. Fonseca, P. H. Souto Ribeiro, S. Pádua, and C. H. Monken, Phys. Rev. A **60**, 1530 (1999);
- [14] B. E. A. Saleh, A. F. Abouraddy, A. V. Sergienko, and M. C. Teich, Phys. Rev. A **62**, 043816 (2000).
- [15] A. F. Abouraddy, B. E. A. Saleh, A. V. Sergienko, and M. C. Teich, Phys. Rev. Lett. **87**, 123602 (2001).
- [16] A. Gatti, E. Brambilla, and L. A. Lugiato, Phys. Rev. Lett. **90**, 133603 (2003).
- [17] R. S. Bennink, S. J. Bentley and R. W. Boyd, Phys. Rev. Lett. **89**, 113601 (2002); R. S. Bennink, S. J. Bentley, R. W. Boyd, J. C. Howell, Phys. Rev. Lett. **92**, 033601 (2004).
- [18] A. Gatti, E. Brambilla, M. Bache and L. A. Lugiato, quant-ph/0307187.
- [19] D. Zh. Cao and K. Wang, quant-ph/0310046; D. Zh. Cao, Zh. Li, Y. H. Zhai, and K. Wang, quant-ph/0401109.


Captions of Figures

Fig. 1 Coincidence interference patterns for the different normalized bandwidths $W = wb/(2\pi)$ of spatial frequency spectrum of a thermal light. $X = xk_0b/(2\pi f)$ is the normalized position in the detection plane and the double slit parameter is taken as $d = 4b$.

Fig. 2 (a) Schematic diagram of coincidence subwavelength interference for a thermal light. (b) The right and left sides show respectively the field $E(q)$ and its conjugate $E^*(q)$ to be correlated, resulting in twice the optical path difference.



a



b

